

# Natural Radioactivity Measurements in Fine Aerosols

Jeff Gaffney, Nancy Marley (University of Arkansas at Little Rock),  
Neil C. Sturchio, and Nancy Martinez (University of Illinois, Chicago)


## ABSTRACT

Samples were collected at the MILAGRO T0 and T1 sites for measurements of  $^7\text{Be}$ ,  $^{210}\text{Pb}$ ,  $^{210}\text{Po}$ , and  $^{210}\text{Bi}$  on 8x10 inch quartz fiber filters at 12 hour increments (day - night) using high volume impactors with 0.1 to 1.0 micron size cuts.  $^7\text{Be}$  and  $^{210}\text{Pb}$  were determined directly by using gamma counting instrumentation. Portions of the samples were analyzed for  $^{210}\text{Po}$  (138day half-life) and  $^{210}\text{Bi}$  (5-day half-life) by using a solid phase extraction method developed at Argonne National Laboratory (1). The samples were dissolved in nitric acid and then treated with HCL to form the metal chloride complexes. The  $^{210}\text{Po}$  and  $^{210}\text{Bi}$  can then be easily separated from the parent  $^{210}\text{Pb}$  by filtering the solution through extraction membranes that have been impregnated with particles containing the strongly basic quarternary amine groups. The filter membranes were then counted to determine the adsorbed  $^{210}\text{Bi}$  and  $^{210}\text{Po}$  using beta and alpha counting equipment.






The samples will also be separated into organic and elemental carbon fractions and analyzed for  $^{14}\text{C}$ ,  $^{13}\text{C}$ , and  $^{12}\text{C}$  using accelerator mass spectrometry and isotope ratio mass spectrometry, respectively. These analysis are still underway and will be reported at a later date.

This work was supported by the DOE Atmospheric Science Program as part of the Megacity Aerosol Experiment - Mexico City (MAX-Mex) portion of MILAGRO. Portions of this work were conducted at Argonne National Laboratory.

1. N.A. Marley, J.S. Gaffney, K.A. Orlandini, P.J. Drayton, and M.M. Cunningham "An improved method for the separation of  $^{210}\text{Bi}$  and  $^{210}\text{Po}$  from  $^{210}\text{Pb}$  by using solid-phase extraction disk membranes: Environmental Applications" *Radiochimica Acta*, 85, 71-78 (1999).





The parent Pb-210 can easily and quickly be separated from the progeny Bi-210 and Po-210 in their chloride forms.



Be-7 values were low during the study period indicating little upper atmospheric mixing. Bi-210 and Po-210 were similar at both sites and preliminary comparison of Po-210/Bi210 ratios indicate an average aerosol residence time of ~30 days at both sites.

Work on Pb-210, K-40, C-13, and C-14 is ongoing. These results will help determine the aerosol atmospheric residence times as well as % biogenic vs % fossil carbon in both organic and elemental aerosol fractions.



## ACKNOWLEDGEMENT

This work was supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Atmospheric Science Program. A portion of this work was done at Argonne National Laboratory under contract W-31-109-Eng-38.